



HSL Phospho-Ser660 rabbit pAb

Catalog No	YP-Ab-10570
Isotype	IgG
Reactivity	Human; Mouse; Rat
Applications	WB
Gene Name	LIPE
Protein Name	HSL Ser660
Immunogen	Synthesized peptide derived from human HSL Ser660
Specificity	This antibody detects endogenous levels of HSL Ser660 at Human, Mouse, Rat
Formulation	Liquid in PBS containing 50% glycerol, and 0.226% sodium azide.
Source	Polyclonal, Rabbit, IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Hormone-sensitive lipase (HSL) (EC 3.1.1.79)
Observed Band	
Cell Pathway	Cell membrane . Membrane, caveola . Cytoplasm, cytosol . Lipid droplet . Found in the high-density caveolae. Translocates to the cytoplasm from the caveolae upon insulin stimulation (PubMed:17026959). Phosphorylation by AMPK reduces its translocation towards the lipid droplets (By similarity).
Tissue Specificity	Testis.
Function	catalytic activity: Diacylglycerol + H ₂ O = monoacylglycerol + a carboxylate., catalytic activity: Monoacylglycerol + H ₂ O = glycerol + a carboxylate., catalytic activity: Triacylglycerol + H ₂ O = diacylglycerol + a carboxylate., enzyme regulation: Rapidly activated by cAMP-dependent phosphorylation under the influence of catecholamines. Dephosphorylation and inactivation are controlled by insulin., function: In adipose tissue and heart, it primarily hydrolyzes stored triglycerides to free fatty acids, while in steroidogenic tissues, it principally converts cholesteryl esters to free cholesterol for steroid hormone production., pathway: Glycerolipid metabolism; triacylglycerol degradation., similarity: Belongs to the 'GDXG' lipolytic enzyme family., subcellular location: Found in the high-density caveolae. Translocates to the cytoplasm from the caveolae upon insulin stimulation., subunit: Interacts wi

Background

The protein encoded by this gene has a long and a short form, generated by use of alternative translational start codons. The long form is expressed in steroidogenic tissues such as testis, where it converts cholesteryl esters to free cholesterol for steroid hormone production. The short form is expressed in adipose tissue, among others, where it hydrolyzes stored triglycerides to free fatty acids. [provided by RefSeq, Jul 2008],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images