



Mac-2BP Polyclonal Antibody

Catalog No	YP-Ab-10642
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	LGALS3BP M2BP
Protein Name	Galectin-3-binding protein (Basement membrane autoantigen p105) (Lectin galactoside-binding soluble 3-binding protein) (Mac-2-binding protein) (MAC2BP) (Mac-2 BP) (Tumor-associated antigen 90K)
Immunogen	Synthetic peptide from human protein at AA range: 100-160
Specificity	The antibody detects endogenous Mac-2BP
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000, ELISA 1:10000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Galectin-3-binding protein (Basement membrane autoantigen p105) (Lectin galactoside-binding soluble 3-binding protein) (Mac-2-binding protein) (MAC2BP) (Mac-2 BP) (Tumor-associated antigen 90K)
Observed Band	65kD
Cell Pathway	Secreted . Secreted, extracellular space, extracellular matrix .
Tissue Specificity	Ubiquitous. Detected in body fluids such as semen, milk, serum, tears, saliva and urine. Expressed by keratinocytes and fibroblasts.
Function	function:Promotes intergrin-mediated cell adhesion. May stimulate host defense against viruses and tumor cells.,similarity:Contains 1 BACK (BTB/Kelch associated) domain.,similarity:Contains 1 BTB (POZ) domain.,similarity:Contains 1 SRCR domain.,subunit:Homodimers and homomultimers. The multimers form ring-like structures with a diameter of 30-40 nm. Binds LGALS1 and LGALS3. Binds ITGB1, COL4A1, COL5A1, COL6A1, FN1 and NID.,tissue specificity:Ubiquitous. Detected in body fluids such as semen, milk, serum, tears, saliva and urine. Expressed by keratinocytes and fibroblasts.,
Background	The galectins are a family of beta-galactoside-binding proteins implicated in modulating cell-cell and cell-matrix interactions. LGALS3BP has been found elevated in the serum of patients with cancer and in those infected by the human immunodeficiency virus (HIV). It appears to be implicated in immune response



associated with natural killer (NK) and lymphokine-activated killer (LAK) cell cytotoxicity. Using fluorescence in situ hybridization the full length 90K cDNA has been localized to chromosome 17q25. The native protein binds specifically to a human macrophage-associated lectin known as Mac-2 and also binds galectin 1. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of human blood lysate, antibody was diluted at 1000. Secondary antibody(catalog#:RS0002) was diluted at 1:20000