

(Tel: 400-999-8863 ■ Emall:Upingbio.163.com



Tetranectin Polyclonal Antibody

| Catalog No | YP-Ab-10709 |
|---------------------------|--|
| Isotype | IgG |
| Reactivity | Human;Mouse |
| Applications | IHC;IF;ELISA |
| Gene Name | CLEC3B TNA |
| Protein Name | Tetranectin (TN) (C-type lectin domain family 3 member B) (Plasminogen kringle 4-binding protein) |
| Immunogen | Synthetic peptide from human protein at AA range: 11-60 |
| Specificity | The antibody detects endogenous Tetranectin |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Polyclonal, Rabbit,IgG |
| Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | IHC-p 1:50-200, ELISA 1:10000-20000. IF 1:50-200 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | Tetranectin (TN;C-type lectin domain family 3 member B;Plasminogen kringle 4-binding protein) |
| Observed Band | |
| Cell Pathway | Secreted. |
| Tissue Specificity | Found in plasma. |
| Function | function:Tetranectin binds to plasminogen and to isolated kringle 4. May be involved in the packaging of molecules destined for exocytosis.,mass spectrometry: PubMed:10614823,online information:Tetranectin,similarity:Contains 1 C-type lectin domain.,subunit:Homotrimer.,tissue specificity:Found in plasma., |
| Background | function:Tetranectin binds to plasminogen and to isolated kringle 4. May be involved in the packaging of molecules destined for exocytosis.,mass spectrometry: PubMed:10614823,online information:Tetranectin,similarity:Contains 1 C-type lectin domain.,subunit:Homotrimer.,tissue specificity:Found in plasma., |
| matters needing attention | Avoid repeated freezing and thawing! |



UpingBio technology Co.,Ltd


(Tel: 400-999-8863 ■ Emall:Upingbio.163.com




Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemical analysis of paraffin-embedded human-kidney, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded human-kidney, antibody was diluted at 1:200