



ASPP1 Polyclonal Antibody

Catalog No	YP-Ab-10728
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	IHC;IF;ELISA
Gene Name	PPP1R13B ASPP1 KIAA0771
Protein Name	Apoptosis-stimulating of p53 protein 1 (Protein phosphatase 1 regulatory subunit 13B)
Immunogen	Synthetic peptide from human protein at AA range: 630-680
Specificity	The antibody detects endogenous ASPP1
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	IHC-p 1:50-200, ELISA 1:10000-20000. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Apoptosis-stimulating of p53 protein 1 (Protein phosphatase 1 regulatory subunit 13B)
Observed Band	
Cell Pathway	Cytoplasm . Nucleus . Predominantly cytoplasmic. Some fraction is nuclear.
Tissue Specificity	Reduced expression in breast carcinomas expressing a wild-type TP53 protein.
Function	disease:Defects in PPP1R13B may be a cause of breast cancers. It is overexpressed in many patients suffering from breast carcinomas and expressing a wild-type TP53 protein.,domain:The ankyrin repeats and the SH3 domain are required for specific interactions with TP53.,function:Regulator that plays a central role in regulation of apoptosis via its interaction with p53/TP53. Regulates TP53 by enhancing the DNA binding and transactivation function of TP53 on the promoters of proapoptotic genes in vivo.,miscellaneous:In contrast to its official gene name, it is not a regulatory subunit of protein phosphatase 1. This name was given due to its similarity with a protein that binds to protein phosphatase 1.,similarity:Belongs to the ASPP family.,similarity:Contains 1 SH3 domain.,similarity:Contains 2 ANK repeats.,subcellular location:Predominantly cytoplasmic. Some fraction is nuclear.,subunit:l
Background	This gene encodes a member of the ASPP (apoptosis-stimulating protein of p53) family of p53 interacting proteins. The protein contains four ankyrin repeats and



an SH3 domain involved in protein-protein interactions. ASPP proteins are required for the induction of apoptosis by p53-family proteins. They promote DNA binding and transactivation of p53-family proteins on the promoters of proapoptotic genes. Expression of this gene is regulated by the E2F transcription factor. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemical analysis of paraffin-embedded Human-liver, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded Human-liver, antibody was diluted at 1:100