

(Tel: 400-999-8863 ■ Email:Upingbio.163.com




TCEAL7 Polyclona Antibody


| Catalog No | YP-Ab-10892 |
|---------------------------|---|
| Isotype | IgG |
| Reactivity | Human; Mouse |
| Applications | WB;IHC;IF;ELISA |
| Gene Name | TCEAL7 |
| Protein Name | TCEAL7 |
| Immunogen | Synthesized peptide derived from human TCEAL7 AA range: 34-84 |
| Specificity | This antibody detects endogenous levels of human TCEAL7 |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Polyclonal, Rabbit,IgG |
| Purification | The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen. |
| Dilution | Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in other applications. |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | Transcription elongation factor A protein-like 7 (TCEA-like protein 7;Transcription elongation factor S-II protein-like 7) |
| Observed Band | |
| Cell Pathway | Nucleus . |
| Tissue Specificity | Highly expressed in normal and fetal brain tissues, and weakly expressed in uterus and ovary. Down-regulated in epithelial ovarian, cervical, prostate, breast, brain and lung cancer cell lines and in brain and ovarian tumors. |
| Function | function:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TFA subfamily., |
| Background | function:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TFA subfamily., |
| matters needing attention | Avoid repeated freezing and thawing! |
| Usage suggestions | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel. |
| | |



Products Images



Western blot analysis of lysates from SH-SY5Y cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human brain tumor. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).