




FCGBP rabbit pAb

Catalog No	YP-Ab-11057
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	IHC;IF
Gene Name	FCGBP
Protein Name	FCGBP
Immunogen	Synthesized peptide derived from human FCGBP AA range: 1806-1856
Specificity	This antibody detects endogenous levels of FCGBP at Human
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	IHC-p 1: 50-200. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Secreted .
Tissue Specificity	Mainly expressed in placenta and colon epithelium. Expressed in thyroid, and down-regulated in thyroid carcinomas. Present in serum, with higher levels in patients with various autoimmune diseases (at protein level).
Function	domain:The N-terminal IgGFc-binding region is primate-specific.,function:May be involved in the maintenance of the mucosal structure as a gel-like component of the mucosa.,similarity:Contains 12 TIL (trypsin inhibitory-like) domains.,similarity:Contains 13 VWFD domains.,subunit:Interacts with the Fc portion of IgG and with MUC2.,tissue specificity:Mainly expressed in placenta and colon epithelium. Expressed in thyroid, and down-regulated in thyroid carcinomas. Present in serum, with higher levels in patients with various autoimmune diseases (at protein level).,
Background	
matters needing attention	Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).