








FA69B rabbit pAb

Catalog No	YP-Ab-11066
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	WB
Gene Name	FAM69B C9orf136 PP6977
Protein Name	FA69B
Immunogen	Synthesized peptide derived from human FA69B AA range: 333-383
Specificity	This antibody detects endogenous levels of FA69B at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	1 mg/ml
Concentration Purity	1 mg/ml ≥90%
	-
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90%
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year Endoplasmic reticulum membrane ; Single-pass type II membrane protein . similarity:Belongs to the FAM69 family., This gene encodes a member of the FAM69 family of cysteine-rich type II
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90% -20°C/1 year Endoplasmic reticulum membrane ; Single-pass type II membrane protein . similarity:Belongs to the FAM69 family., This gene encodes a member of the FAM69 family of cysteine-rich type II transmembrane proteins. These proteins localize to the endoplasmic reticulum but
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	≥90% -20°C/1 year Endoplasmic reticulum membrane ; Single-pass type II membrane protein . similarity:Belongs to the FAM69 family., This gene encodes a member of the FAM69 family of cysteine-rich type II transmembrane proteins. These proteins localize to the endoplasmic reticulum but their specific functions are unknown. [provided by RefSeq, Nov 2011],



Products Images



Western blot analysis of lysates from VEC cells, primary antibody was diluted at 1:1000, 4° over night