

🔇 Tel: 400-999-8863 💌 Email:UpingBio@163.com

Website: www.upingBio.com

## TLCD1 rabbit pAb


Catalog No	YP-Ab-11123
Isotype	lgG
Reactivity	Human; Mouse;Rat
Applications	WB;ELISA;IHC
Gene Name	TLCD1
Protein Name	TLCD1
Immunogen	Synthesized peptide derived from human TLCD1 AA range: 162-212
Specificity	This antibody detects endogenous levels of TLCD1 at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000
0	1 mg/ml
Concentration	T mg/mi
Purity	≥90%
	-
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90%
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year Cell membrane ; Multi-pass membrane protein .
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90% -20°C/1 year Cell membrane ; Multi-pass membrane protein .
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	≥90% -20°C/1 year Cell membrane ; Multi-pass membrane protein . similarity:Contains 1 TLC (TRAM/LAG1/CLN8) domain.,




🔇 Tel: 400-999-8863 💌 Email:UpingBio@163.com

Ø Website: www.upingBio.com

## **Products Images**



Western blot analysis of lysates from SW480 cells, primary antibody was diluted at 1:1000, 4°over night



Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).