





ATD3A rabbit pAb

YP-Ab-11167
IgG
Human;Rat;Mouse;
WB
ATAD3A
ATD3A
Synthesized peptide derived from human ATD3A AA range: 433-483
This antibody detects endogenous levels of ATD3A at Human
Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Polyclonal, Rabbit,IgG
The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
WB 1: 500-2000
1 mg/ml
≥90%
-20°C/1 year
Mitochondrion inner membrane; Single-pass membrane protein. Mitochondrion matrix, mitochondrion nucleoid. In the mitochondrial inner membrane, enriched in sites with the potential to form contacts with the outer membrane (PubMed:20349121, PubMed:20154147). The N-terminal domain interacts with the inner surface of the mitochondrial outer membrane and the C-terminal domain localizes in a specific matrix compartment, where it is associated with nucleoids (PubMed:18063578).
Overexpressed in lung adenocarcinomas (at protein level).
similarity:Belongs to the AAA ATPase family.,
, , ,
ATAD3A and ATAD3B (MIM 612317) are mitochondrial membrane proteins that contribute to the stabilization of large mitochondrial DNA (mtDNA)-protein complexes called nucleoids (He et al., 2007 [PubMed 17210950]).[supplied by OMIM, Sep 2008],



UpingBio technology Co.,Ltd


C Tel: 400-999-8863 ■ Email:UpingBio@163.com



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from HpeG2 cells, primary antibody was diluted at 1:1000, 4° over night