







# CAR16 rabbit pAb

Catalog No	YP-Ab-11168
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	CARD16 COP COP1
Protein Name	CAR16
Immunogen	Synthesized peptide derived from human CAR16 AA range: 101-151
Specificity	This antibody detects endogenous levels of CAR16 at Human
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Storage Stability Synonyms	-20°C/1 year
	-20°C/1 year

Widely expressed. Expressed at higher level in placenta, spleen, lymph node and Tissue Specificity bone marrow. Weakly or not expressed in thymus.

#### **Function**

function: Caspase inhibitor. Acts as a regulator of procaspase-1/CASP1 activation implicated in the regulation of the proteolytic maturation of pro-interleukin-1 beta (IL1B) and its release during inflammation. Inhibits the release of IL1B in response to LPS in monocytes. Also induces NF-kappa-B activation during the pro-inflammatory cytokine response. Also able to inhibit CASP1-mediated neuronal cell death, TNF-alpha, hypoxia-, UV-, and staurosporine-mediated cell death but not ER stress-mediated cell death. Acts by preventing activation of caspases CASP1 and CASP4, possibly by preventing the interaction between CASP1 and RIPK2.,induction:Down-regulated in patients suffering of Huntington disease.,similarity:Contains 1 CARD domain.,subunit:Homooligomer. Interacts with CASP1, CASP4, CARD8 and RIPK2.,tissue specificity:Widely expressed. Expressed at higher level in placenta, spleen, lymph Expressed at higher level in placenta, spleen, lymph

#### **Background**



### UpingBio technology Co.,Ltd








matters needing attention

Avoid repeated freezing and thawing!

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western blot analysis of lysates from Hela cells, primary antibody was diluted at 1:1000, 4°over night