








## WDR54 rabbit pAb

Catalog No	YP-Ab-11191
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB
Gene Name	WDR54
Protein Name	WDR54
Immunogen	Synthesized peptide derived from human WDR54 AA range: 99-149
Specificity	This antibody detects endogenous levels of WDR54 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	1 mg/ml
<b>Concentration Purity</b>	1 mg/ml ≥90%
	•
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90%
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year  Vesicle . Aggregates in vesicles when cross-linked  Expressed in epithelial cells (at protein level) (PubMed:29987896). Isoform 3
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90%  -20°C/1 year  Vesicle . Aggregates in vesicles when cross-linked  Expressed in epithelial cells (at protein level) (PubMed:29987896). Isoform 3 expression is highly increased in colorectal cancer cells (PubMed:29987896).
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90%  -20°C/1 year  Vesicle . Aggregates in vesicles when cross-linked  Expressed in epithelial cells (at protein level) (PubMed:29987896). Isoform 3 expression is highly increased in colorectal cancer cells (PubMed:29987896).
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	≥90%  -20°C/1 year  Vesicle . Aggregates in vesicles when cross-linked  Expressed in epithelial cells (at protein level) (PubMed:29987896). Isoform 3 expression is highly increased in colorectal cancer cells (PubMed:29987896). similarity:Contains 4 WD repeats.,



## **Products Images**



Western blot analysis of lysates from HEK293 cells, primary antibody was diluted at 1:1000, 4° over night