






SHSA3 rabbit pAb


Catalog No	YP-Ab-11261
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB;ELISA;IHC
Gene Name	SHISA3
Protein Name	SHSA3
Immunogen	Synthesized peptide derived from human SHSA3 AA range: 126-176
Specificity	This antibody detects endogenous levels of SHSA3 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000
Concentration	1 mg/ml
	-
Purity	≥90%
Purity Storage Stability	
	≥90%
Storage Stability	≥90%
Storage Stability Synonyms	≥90%
Storage Stability Synonyms Observed Band	≥90% -20°C/1 year
Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year Endoplasmic reticulum membrane ; Single-pass type I membrane protein . function:Plays an essential role in the maturation of presomitic mesoderm cells by individual attenuation of both FGF and WNT signaling.,similarity:Belongs to the
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90% -20°C/1 year Endoplasmic reticulum membrane ; Single-pass type I membrane protein . function:Plays an essential role in the maturation of presomitic mesoderm cells by individual attenuation of both FGF and WNT signaling.,similarity:Belongs to the
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	≥90% -20°C/1 year Endoplasmic reticulum membrane ; Single-pass type I membrane protein . function:Plays an essential role in the maturation of presomitic mesoderm cells by individual attenuation of both FGF and WNT signaling.,similarity:Belongs to the shisa protein family.,



Products Images



Western blot analysis of lysates from HpeG2 cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).