

🔇 Tel: 400-999-8863 💌 Email:UpingBio@163.com



LRRN3 rabbit pAb

| Catalog No | YP-Ab-11292 |
|---|--|
| Isotype | IgG |
| Reactivity | Human; Mouse;Rat |
| Applications | WB;IHC |
| Gene Name | LRRN3 Nbla10363 UNQ194/PRO220 |
| Protein Name | LRRN3 |
| Immunogen | Synthesized peptide derived from human LRRN3 AA range: 225-275 |
| Specificity | This antibody detects endogenous levels of LRRN3 at Human/Mouse/Rat |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Polyclonal, Rabbit,IgG |
| Purification | The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. |
| Dilution | WB 1:500-2000;IHC-p 1:50-300 |
| Concentration | 1 mg/ml |
| | |
| Purity | ≥90% |
| Purity Storage Stability | ≥90% -20°C/1 year |
| - | |
| Storage Stability | |
| Storage Stability Synonyms | |
| Storage Stability Synonyms Observed Band | -20°C/1 year |
| Storage Stability Synonyms Observed Band Cell Pathway | -20°C/1 year |
| Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity | -20°C/1 year Membrane ; Single-pass type I membrane protein . similarity:Contains 1 fibronectin type-III domain.,similarity:Contains 1 Ig-like C2-type (immunoglobulin-like) domain.,similarity:Contains 12 LRR (leucine-rich) |
| Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function | -20°C/1 year Membrane ; Single-pass type I membrane protein . similarity:Contains 1 fibronectin type-III domain.,similarity:Contains 1 Ig-like C2-type (immunoglobulin-like) domain.,similarity:Contains 12 LRR (leucine-rich) |
| Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing | -20°C/1 year Membrane ; Single-pass type I membrane protein . similarity:Contains 1 fibronectin type-III domain.,similarity:Contains 1 Ig-like C2-type (immunoglobulin-like) domain.,similarity:Contains 12 LRR (leucine-rich) repeats., |




🕓 Tel: 400-999-8863 💌 Email:UpingBio@163.com

Ø Website: www.upingBio.com

w.upingBio.com

Products Images



Lin hi di di

Western blot analysis of lysates from Jarkat cells, primary antibody was diluted at 1:1000, 4°over night

Immunohistochemical analysis of paraffin-embedded human Gastric adenocarcinoma. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).