







ARFRP rabbit pAb

Catalog No	YP-Ab-11323
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	WB
Gene Name	ARFRP1 ARP1
Protein Name	ARFRP
Immunogen	Synthesized peptide derived from human ARFRP AA range: 110-160
Specificity	This antibody detects endogenous levels of ARFRP at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Call Dethyray	
Cell Pathway	Golgi apparatus . Golgi apparatus, trans-Golgi network . Located in the trans-Golgi in the GTP-bound active state
Tissue Specificity	Golgi apparatus . Golgi apparatus, trans-Golgi network . Located in the trans-Golgi in the GTP-bound active state Found in most tissues.
-	trans-Golgi in the GTP-bound active state
Tissue Specificity	trans-Golgi in the GTP-bound active state Found in most tissues. function:Possibly involved in plasma membrane-related signaling events.,similarity:Belongs to the small GTPase superfamily. Arf
Tissue Specificity Function	trans-Golgi in the GTP-bound active state Found in most tissues. function:Possibly involved in plasma membrane-related signaling events.,similarity:Belongs to the small GTPase superfamily. Arf family.,subunit:Interacts with SYS1.,tissue specificity:Found in most tissues., The protein encoded by this gene is a membrane-associated GTP-ase which localizes to the plasma membrane and is related to the ADP-ribosylation factor (ARF) and ARF-like (ARL) proteins. This gene plays a role in membrane trafficking between the trans-Golgi network and endosomes. Alternatively spliced transcript variants encoding different isoforms have been identified. [provided by



UpingBio technology Co.,Ltd


C Tel: 400-999-8863 ≤ Email:UpingBio@163.com



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from Hela cells, primary antibody was diluted at 1:1000, 4° over night