



NEUR3 rabbit pAb

Catalog No	YP-Ab-11370
Isotype	IgG
Reactivity	Human; Mouse; Rat
Applications	WB
Gene Name	NEU3
Protein Name	NEUR3
Immunogen	Synthesized peptide derived from human NEUR3 AA range: 132-182
Specificity	This antibody detects endogenous levels of NEUR3 at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit, IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Cell membrane ; Peripheral membrane protein . Membrane, caveola . Early endosome membrane ; Peripheral membrane protein . Recycling endosome membrane ; Peripheral membrane protein . Lysosome membrane ; Peripheral membrane protein . Associates with the external leaflet of the plasma membrane (By similarity). S-acylated NEU3 likely spans the lipid bilayer with a portion of C-terminus exposed to the cytosol and the catalytic region facing the extracellular space (PubMed:28646141). .
Tissue Specificity	Highly expressed in skeletal muscle, testis, adrenal gland and thymus, followed by pancreas, liver, heart and thymus. Weakly expressed in kidney, placenta, brain and lung.
Function	catalytic activity:Hydrolysis of alpha-(2->3)-, alpha-(2->6)-, alpha-(2->8)-glycosidic linkages of terminal sialic acid residues in oligosaccharides, glycoproteins, glycolipids, colominic acid and synthetic substrates.,function:Plays a role in modulating the ganglioside content of the lipid bilayer at the level of membrane-bound sialyl glycoconjugates.,similarity:Belongs to the glycosyl hydrolase 33 family.,similarity:Contains 3 BNR repeats.,tissue specificity:Highly expressed in skeletal muscle, testis, adrenal gland and thymus, followed by pancreas, liver, heart and thymus. Weakly expressed in kidney, placenta, brain and lung.,



Background

This gene product belongs to a family of glycohydrolytic enzymes which remove sialic acid residues from glycoproteins and glycolipids. It is localized in the plasma membrane, and its activity is specific for gangliosides. It may play a role in modulating the ganglioside content of the lipid bilayer. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from A431 cells, primary antibody was diluted at 1:1000, 4° over night