


## NCEH1 rabbit pAb


Catalog No	YP-Ab-11477
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	WB;ELISA;IHC
Gene Name	NCEH1 AADACL1 KIAA1363
Protein Name	NCEH1
Immunogen	Synthesized peptide derived from human NCEH1 AA range: 8-58
Specificity	This antibody detects endogenous levels of NCEH1 at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000
Concentration	1 mg/ml
Concentration Purity	1 mg/ml ≥90%
	•
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90%
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90%  -20°C/1 year  Cell membrane ; Single-pass type II membrane protein . Microsome .  Expressed in monocyte-derived macrophages. Up-regulated in invasive
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90%  -20°C/1 year  Cell membrane ; Single-pass type II membrane protein . Microsome .  Expressed in monocyte-derived macrophages. Up-regulated in invasive
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90%  -20°C/1 year  Cell membrane ; Single-pass type II membrane protein . Microsome .  Expressed in monocyte-derived macrophages. Up-regulated in invasive
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	≥90%  -20°C/1 year  Cell membrane ; Single-pass type II membrane protein . Microsome .  Expressed in monocyte-derived macrophages. Up-regulated in invasive melanoma and breast carcinoma cell lines.



## **Products Images**



Western blot analysis of lysates from MCF-7 cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).