





EIF3I rabbit pAb

Catalog No	YP-Ab-11506
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	WB
Gene Name	EIF3I EIF3S2 TRIP1
Protein Name	EIF3I
Immunogen	Synthesized peptide derived from human EIF3I AA range: 100-150
Specificity	This antibody detects endogenous levels of EIF3I at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Cytoplasm .
Tissue Specificity	
Function	function:Component of the eukarvotic translation initiation factor 3 (eIF-3)

function: Component of the eukaryotic translation initiation factor 3 (eIF-3) Function

function:Component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis. The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNAi and eIF-5 to form the 43S preinitiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of posttermination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation.,mass spectrometry: PubMed:17322308,mass spectrometry: PubMed:18599441,PTM:Phosphorylated by TGF-beta type II receptor.,similarity:Belongs to the eIF-3 subunit I family..similarity:Contains 5 WD repeats..subunit:Component of the eukaryotic

family, similarity: Contains 5 WD repeats, subunit: Component of the eukaryotic

Background



UpingBio technology Co.,Ltd

€ Tel: 400-999-8863 🛎 Email:UpingBio@163.com




matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from HpeG2 cells, primary antibody was diluted at 1:1000, 4°over night