





## PCLI1 rabbit pAb

Catalog No	YP-Ab-11574
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB;ELISA;IHC
Gene Name	PID1 NYGGF4 PCLI1 HMFN2073
Protein Name	PCLI1
Immunogen	Synthesized peptide derived from human PCLI1 AA range: 144-194
Specificity	This antibody detects endogenous levels of PCLI1 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Cytoplasm .
Tissue Specificity	Expressed in subcutaneous fat, heart, skeletal muscle, brain, colon, thymus, spleen, kidney, liver, small intestine, placenta, lung and peripheral blood leukocyte.
Function	function:Increases proliferation of preadipocytes without affecting adipocytic differentiation.,induction:Up-regulated in fat of obese subjects.,polymorphism:Some sequences seem to have a duplication of exon 2.,similarity:Contains 1 PID domain.,subunit:Found in a complex with PID1/PCLI1 LRP1 and CUBNI. Interacts with LRP1 and CUBN.,tissue specificity:Expressed in subcutaneous fat, heart, skeletal muscle, brain, colon, thymus, spleen, kidney, liver, small intestine, placenta, lung and peripheral blood leukocyte.,
Background	
matters needing attention	Avoid repeated freezing and thawing!



## UpingBio technology Co.,Ltd


C Tel: 400-999-8863 ■ Email:UpingBio@163.com




**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western blot analysis of lysates from Hela cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).