







TBCD7 rabbit pAb

Catalog No	YP-Ab-11605
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB
Gene Name	TBC1D7 HSPC239 TBC7
Protein Name	TBCD7
Immunogen	Synthesized peptide derived from human TBCD7 AA range: 85-135
Specificity	This antibody detects endogenous levels of TBCD7 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Cytoplasmic vesicle . Localizes in the cytoplasmic vesicles of the endomembrane in association with TSC1-TSC2 complex.
	Cytoplasmic vesicle . Localizes in the cytoplasmic vesicles of the endomembrane in association with TSC1-TSC2 complex. Highly expressed in heart, and slightly in kidney, liver and placenta.
Cell Pathway	in association with TSC1-TSC2 complex.
Cell Pathway Tissue Specificity	in association with TSC1-TSC2 complex. Highly expressed in heart, and slightly in kidney, liver and placenta. function:May act as a GTPase-activating protein for Rab family protein(s)
Cell Pathway Tissue Specificity Function	Highly expressed in heart, and slightly in kidney, liver and placenta. function:May act as a GTPase-activating protein for Rab family protein(s) .,similarity:Contains 1 Rab-GAP TBC domain., This gene encodes a member of the TBC-domain containing protein family. The encoded protein functions as a subunit of the tuberous sclerosis TSC1-TSC2 complex which plays a role in the regulation of cellular growth and differentiation. Mutations in this gene have been associated with autosomal recessive megalencephaly. Alternative splicing results in multiple transcript variants. Naturally occurring readthrough transcription occurs between this locus and



UpingBio technology Co.,Ltd


C Tel: 400-999-8863 ■ Email:UpingBio@163.com



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from DU145 cells, primary antibody was diluted at 1:1000, 4° over night