








PKHA7 rabbit pAb

Catalog No	YP-Ab-11629
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB;ELISA;IHC
Gene Name	PLEKHA7
Protein Name	PKHA7
Immunogen	Synthesized peptide derived from human PKHA7 AA range: 96-146
Specificity	This antibody detects endogenous levels of PKHA7 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Cell junction, adherens junction. Cytoplasm. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Localizes to zonula adherens, recruited via its interaction with CTNND1
Cell Pathway Tissue Specificity	Cell junction, adherens junction . Cytoplasm . Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Localizes to zonula adherens, recruited via its interaction with CTNND1
•	Cell junction, adherens junction. Cytoplasm. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Localizes to zonula adherens, recruited via its interaction with CTNND1
Tissue Specificity	microtubule organizing center, centrosome . Localizes to zonula adherens, recruited via its interaction with CTNND1
Tissue Specificity Function	microtubule organizing center, centrosome . Localizes to zonula adherens, recruited via its interaction with CTNND1
Tissue Specificity Function Background matters needing	microtubule organizing center, centrosome . Localizes to zonula adherens, recruited via its interaction with CTNND1 similarity:Contains 1 PH domain.,similarity:Contains 2 WW domains.,






Products Images



Western blot analysis of lysates from PC-12 cells, primary antibody was diluted at 1:1000, 4°over night



Immunohistochemical analysis of paraffin-embedded human Squamous cell carcinoma of lung. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).