








TRI44 rabbit pAb

Catalog No	YP-Ab-11694
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	WB
Gene Name	TRIM44 DIPB
Protein Name	TRI44
Immunogen	Synthesized peptide derived from human TRI44 AA range: 42-92
Specificity	This antibody detects endogenous levels of TRI44 at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	1 mg/ml
Purity	≥90%
•	
Storage Stability	-20°C/1 year
Storage Stability Synonyms	-20°C/1 year
	-20°C/1 year
Synonyms	-20°C/1 year intracellular,
Synonyms Observed Band	
Synonyms Observed Band Cell Pathway	intracellular,
Synonyms Observed Band Cell Pathway Tissue Specificity	intracellular, Highly expressed in testis. function:May play a role in the process of differentiation and maturation of neuronal cells.,similarity:Contains 1 B box-type zinc finger., This gene encodes a member of the tripartite motif (TRIM) family. The TRIM motif
Synonyms Observed Band Cell Pathway Tissue Specificity Function	intracellular, Highly expressed in testis. function:May play a role in the process of differentiation and maturation of neuronal cells.,similarity:Contains 1 B box-type zinc finger., This gene encodes a member of the tripartite motif (TRIM) family. The TRIM motif includes three zinc-binding domains, namely a RING, a B-box type 1 and a B-box
Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	intracellular, Highly expressed in testis. function:May play a role in the process of differentiation and maturation of neuronal cells.,similarity:Contains 1 B box-type zinc finger., This gene encodes a member of the tripartite motif (TRIM) family. The TRIM motif includes three zinc-binding domains, namely a RING, a B-box type 1 and a B-box type 2, and a coiled-coil region. [provided by RefSeq, Jul 2008],



Products Images



Western blot analysis of lysates from Hela cells, primary antibody was diluted at 1:1000, 4° over night