

LRC48 rabbit pAb

Catalog No	YP-Ab-11702
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	WB;IHC
Gene Name	LRRC48
Protein Name	LRC48
Immunogen	Synthesized peptide derived from human LRC48 AA range: 250-300
Specificity	This antibody detects endogenous levels of LRC48 at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300
Concentration	1 mg/ml
Concentration Purity	1 mg/ml ≥90%
	-
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90%
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year Cytoplasm, cytoskeleton, cilium axoneme . Cell projection, cilium . Cytoplasm,
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year Cytoplasm, cytoskeleton, cilium axoneme . Cell projection, cilium . Cytoplasm,
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year Cytoplasm, cytoskeleton, cilium axoneme . Cell projection, cilium . Cytoplasm, cytoskeleton, flagellum axoneme . Cell projection, cilium, flagellum .
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90% -20°C/1 year Cytoplasm, cytoskeleton, cilium axoneme . Cell projection, cilium . Cytoplasm, cytoskeleton, flagellum axoneme . Cell projection, cilium, flagellum .
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	≥90% -20°C/1 year Cytoplasm, cytoskeleton, cilium axoneme . Cell projection, cilium . Cytoplasm, cytoskeleton, flagellum axoneme . Cell projection, cilium, flagellum . similarity:Contains 5 LRR (leucine-rich) repeats.,



UpingBio technology Co.,Ltd

C Tel: 400-999-8863 ≤ Email:UpingBio@163.com



Products Images Western blot analysis of lysates from Hela cells, primary antibody was diluted at 1:1000, 4°over night

Immunohistochemical analysis of paraffin-embedded human Gastric adenocarcinoma. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).