







KCTD5 rabbit pAb

Catalog No	YP-Ab-11743
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	WB;ELISA;IHC
Gene Name	KCTD5
Protein Name	KCTD5
Immunogen	Synthesized peptide derived from human KCTD5 AA range: 170-220
Specificity	This antibody detects endogenous levels of KCTD5 at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	000014
	-20°C/1 year
Synonyms	-20°C/1 year
Synonyms Observed Band	-20°C/1 year
	Cytoplasm, cytosol . Cytoplasm . Nucleus. Predominantly cytoplasmic, translocated to the nucleus upon interaction with Rep proteins
Observed Band	Cytoplasm, cytosol . Cytoplasm . Nucleus. Predominantly cytoplasmic,
Observed Band Cell Pathway	Cytoplasm, cytosol . Cytoplasm . Nucleus. Predominantly cytoplasmic,
Observed Band Cell Pathway Tissue Specificity	Cytoplasm, cytosol . Cytoplasm . Nucleus. Predominantly cytoplasmic, translocated to the nucleus upon interaction with Rep proteins similarity:Contains 1 BTB (POZ) domain.,subcellular location:Predominantly cytoplasmic, translocated to the nucleus upon interaction with Rep
Observed Band Cell Pathway Tissue Specificity Function	Cytoplasm, cytosol . Cytoplasm . Nucleus. Predominantly cytoplasmic, translocated to the nucleus upon interaction with Rep proteins similarity:Contains 1 BTB (POZ) domain.,subcellular location:Predominantly cytoplasmic, translocated to the nucleus upon interaction with Rep
Observed Band Cell Pathway Tissue Specificity Function Background matters needing	Cytoplasm, cytosol . Cytoplasm . Nucleus. Predominantly cytoplasmic, translocated to the nucleus upon interaction with Rep proteins similarity:Contains 1 BTB (POZ) domain.,subcellular location:Predominantly cytoplasmic, translocated to the nucleus upon interaction with Rep proteins.,subunit:Interacts with adeno-associated virus 2 REP proteins.,



UpingBio technology Co.,Ltd

C Tel: 400-999-8863 ■ Email:UpingBio@163.com



Western blot analysis of lysates from DU145 cells, primary antibody was diluted at 1:1000, 4°over night Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).