







## K1551 rabbit pAb

Catalog No	YP-Ab-11760
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB;IHC
Gene Name	KIAA1551 C12orf35
Protein Name	K1551
Immunogen	Synthesized peptide derived from human K1551 AA range: 262-312
Specificity	This antibody detects endogenous levels of K1551 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300
Concentration	1 mg/ml
Purity	≥90%
Purity Storage Stability	≥90% -20°C/1 year
Storage Stability	
Storage Stability Synonyms	
Storage Stability Synonyms Observed Band	-20°C/1 year
Storage Stability Synonyms Observed Band Cell Pathway	-20°C/1 year
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	-20°C/1 year  Nucleus . Localizes around gamma-tubulin during M phase  sequence caution:Contaminating sequence.,sequence caution:Contaminating
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	-20°C/1 year  Nucleus . Localizes around gamma-tubulin during M phase  sequence caution:Contaminating sequence.,sequence caution:Contaminating
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	-20°C/1 year  Nucleus . Localizes around gamma-tubulin during M phase  sequence caution:Contaminating sequence.,sequence caution:Contaminating sequence. Potential poly-A sequence.,



## UpingBio technology Co.,Ltd

C Tel: 400-999-8863 ■ Email:UpingBio@163.com



## Western blot analysis of lysates from VEC cells, primary antibody was diluted at 1:1000, 4° over night Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).