

MYG1 rabbit pAb

Catalog No	YP-Ab-11768
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	WB; ELISA;IHC
Gene Name	C12orf10
Protein Name	MYG1
Immunogen	Synthesized peptide derived from human MYG1 AA range: 296-346
Specificity	This antibody detects endogenous levels of MYG1 at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000
Concentration	1 mg/ml
Concentration	1 mg/ml
Purity	≥90%
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90%
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year Nucleus, nucleoplasm . Mitochondrion matrix . Nucleus, nucleolus .
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year Nucleus, nucleoplasm . Mitochondrion matrix . Nucleus, nucleolus . Ubiquitously expressed, with highest levels in testis. similarity:Belongs to the UPF0160 (MYG1) familytissue specificity:Ubiquitously
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90% -20°C/1 year Nucleus, nucleoplasm . Mitochondrion matrix . Nucleus, nucleolus . Ubiquitously expressed, with highest levels in testis. similarity:Belongs to the UPF0160 (MYG1) familytissue specificity:Ubiquitously
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	≥90% -20°C/1 year Nucleus, nucleoplasm . Mitochondrion matrix . Nucleus, nucleolus . Ubiquitously expressed, with highest levels in testis. similarity:Belongs to the UPF0160 (MYG1) family.,tissue specificity:Ubiquitously expressed, with highest levels in testis.,



UpingBio technology Co.,Ltd

C Tel: 400-999-8863 ■ Email:UpingBio@163.com



Western blot analysis of lysates from PC-12 cells, primary antibody was diluted at 1:1000, 4°over night Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4°overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).