


VWA8 rabbit pAb

Catalog No	YP-Ab-11839
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB
Gene Name	VWA8 KIAA0564
Protein Name	VWA8
Immunogen	Synthesized peptide derived from human VWA8 AA range: 1853-1903
Specificity	This antibody detects endogenous levels of VWA8 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	1 mg/ml
Concentration Purity	1 mg/ml ≥90%
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90%
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year [Isoform 1]: Mitochondrion .; [Isoform 2]: Mitochondrion . Peroxisome . Localizes
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year [Isoform 1]: Mitochondrion .; [Isoform 2]: Mitochondrion . Peroxisome . Localizes
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year [Isoform 1]: Mitochondrion .; [Isoform 2]: Mitochondrion . Peroxisome . Localizes to peroxisomes in a PEX7-dependent manner
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90% -20°C/1 year [Isoform 1]: Mitochondrion .; [Isoform 2]: Mitochondrion . Peroxisome . Localizes to peroxisomes in a PEX7-dependent manner
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	≥90% -20°C/1 year [Isoform 1]: Mitochondrion .; [Isoform 2]: Mitochondrion . Peroxisome . Localizes to peroxisomes in a PEX7-dependent manner similarity:Contains 1 VWFA domain.,



Products Images



Western blot analysis of lysates from A431 cells, primary antibody was diluted at 1:1000, 4°over night