

VGLL1 rabbit pAb

Catalog No	YP-Ab-11877
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB
Gene Name	VGLL1 TDU
Protein Name	VGLL1
Immunogen	Synthesized peptide derived from human VGLL1 AA range: 84-134
Specificity	This antibody detects endogenous levels of VGLL1 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	
•	Nucleus.
Tissue Specificity	Nucleus.
•	Nucleus. function:May act as a specific coactivator for the mammalian TEFs.,similarity:Belongs to the vestigial family.,subunit:Interacts with TEFs.,
Tissue Specificity	function:May act as a specific coactivator for the mammalian TEFs.,similarity:Belongs to the vestigial family.,subunit:Interacts with TEFs., The protein encoded by this gene binds proteins of the TEA domain family of
Tissue Specificity Function	function:May act as a specific coactivator for the mammalian TEFs.,similarity:Belongs to the vestigial family.,subunit:Interacts with TEFs., The protein encoded by this gene binds proteins of the TEA domain family of transcription factors (TEFs) through the Vg (vestigial) homology region found in its N-terminus. It may thus function as a specific coactivator for the mammalian
Tissue Specificity Function Background matters needing	function:May act as a specific coactivator for the mammalian TEFs.,similarity:Belongs to the vestigial family.,subunit:Interacts with TEFs., The protein encoded by this gene binds proteins of the TEA domain family of transcription factors (TEFs) through the Vg (vestigial) homology region found in its N-terminus. It may thus function as a specific coactivator for the mammalian TEFs. [provided by RefSeq, Sep 2009],










Products Images



Western blot analysis of lysates from Hela cells, primary antibody was diluted at 1:1000, 4° over night