







AMELY rabbit pAb

YP-Ab-11902
IgG
Human
WB
AMELY AMGL AMGY
AMELY
Synthesized peptide derived from human AMELY.AA range 17-57
This antibody detects endogenous levels of AMELY at Human
Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Polyclonal, Rabbit,IgG
The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
WB 1: 500-2000
1 mg/ml
≥90%
-20°C/1 year
Secreted, extracellular space, extracellular matrix.
developmental stage:Transiently but abundantly expressed by ameloblasts during tooth development. Amelogenin is the predominant protein in developing dental enamel.,function:Plays a role in biomineralization. Seems to regulate the formation of crystallites during the secretory stage of tooth enamel development. Thought to play a major role in the structural organization and mineralization of developing enamel.,miscellaneous:This isoform is encoded on the Y chromosome.,similarity:Belongs to the amelogenin family.,
This gene encodes a member of the amelogenin family of extracellular matrix proteins. Amelogenins are involved in biomineralization during tooth enamel development. Mutations in a related gene on chromosome X cause X-linked amelogenesis imperfecta. [provided by RefSeq, Jul 2008],
Avoid repeated freezing and thawing!



UpingBio technology Co.,Ltd


C Tel: 400-999-8863 ■ Email:UpingBio@163.com



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from HuvEc cells, primary antibody was diluted at 1:1000, 4°over night