








## IN80E rabbit pAb

| Catalog No   | YP-Ab-11929   |
|--|---|
| Isotype  | IgG   |
| Reactivity   | Human;Rat   |
| Applications   | WB  |
| Gene Name  | INO80E CCDC95   |
| Protein Name   | IN80E   |
| Immunogen  | Synthesized peptide derived from human IN80E AA range: 45-95  |
| Specificity  | This antibody detects endogenous levels of IN80E at Human/Rat   |
| Formulation  | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.   |
| Source   | Polyclonal, Rabbit,IgG  |
| Purification   | The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.   |
| Dilution   | WB 1: 500-2000  |
| Concentration  | 1 mg/ml   |
| Purity   | ≥90%  |
|  |   |
| Storage Stability  | -20°C/1 year  |
| Storage Stability Synonyms   | -20°C/1 year  |
|  | -20°C/1 year  |
| Synonyms   | -20°C/1 year  Nucleus .   |
| Synonyms Observed Band   |   |
| Synonyms Observed Band Cell Pathway  |   |
| Synonyms  Observed Band  Cell Pathway  Tissue Specificity                                  | Nucleus .  Subunit:Component of the chromatin-remodeling INO80 complex, at least composed of ACTL6A, ACTR5, ACTR8, RVBL1, RVBL2, INO80, INO80B,                             |
| Synonyms  Observed Band  Cell Pathway  Tissue Specificity  Function                        | Nucleus .  Subunit:Component of the chromatin-remodeling INO80 complex, at least composed of ACTL6A, ACTR5, ACTR8, RVBL1, RVBL2, INO80, INO80B,                             |
| Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing | Nucleus .  subunit:Component of the chromatin-remodeling INO80 complex, at least composed of ACTL6A, ACTR5, ACTR8, RVBL1, RVBL2, INO80, INO80B, INO80C, INO80D and INO80E., |



## **Products Images**



Western blot analysis of lysates from U2OS cells, primary antibody was diluted at 1:1000, 4°over night