







FA69C rabbit pAb

Catalog No	YP-Ab-11941
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB
Gene Name	FAM69C C18orf51
Protein Name	FA69C
Immunogen	Synthesized peptide derived from human FA69C AA range: 345-395
Specificity	This antibody detects endogenous levels of FA69C at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	1 mg/ml
Purity Purity	1 mg/ml ≥90%
	•
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90%
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year Endoplasmic reticulum membrane ; Single-pass type II membrane protein . This gene encodes a member of the FAM69 family of cysteine-rich type II
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90% -20°C/1 year Endoplasmic reticulum membrane ; Single-pass type II membrane protein . This gene encodes a member of the FAM69 family of cysteine-rich type II transmembrane proteins. These proteins localize to the endoplasmic reticulum but
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	≥90% -20°C/1 year Endoplasmic reticulum membrane ; Single-pass type II membrane protein . This gene encodes a member of the FAM69 family of cysteine-rich type II transmembrane proteins. These proteins localize to the endoplasmic reticulum but their specific functions are unknown. [provided by RefSeq, Nov 2011],




UpingBio technology Co.,Ltd





Products Images



Western blot analysis of lysates from U2OS cells, primary antibody was diluted at 1:1000, 4° over night