







SAHH3 rabbit pAb

Catalog No	YP-Ab-11969
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB
Gene Name	AHCYL2 KIAA0828
Protein Name	SAHH3
Immunogen	Synthesized peptide derived from human SAHH3 AA range: 157-207
Specificity	This antibody detects endogenous levels of SAHH3 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	1 mg/ml
Concentration Purity	1 mg/ml ≥90%
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90%
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year Cytoplasm . Microsome . Associates with membranes when phosphorylated,
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year Cytoplasm . Microsome . Associates with membranes when phosphorylated,
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year Cytoplasm . Microsome . Associates with membranes when phosphorylated, probably through interaction with ITPR1 catalytic activity:S-adenosyl-L-homocysteine + H(2)O = L-homocysteine + adenosine.,cofactor:Binds 1 NAD per subunit.,cofactor:NAD.,pathway:Amino-acid biosynthesis; homocysteine biosynthesis; L-homocysteine from S-adenosyl-L-homocysteine: step 1/1.,similarity:Belongs to the
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90% -20°C/1 year Cytoplasm . Microsome . Associates with membranes when phosphorylated, probably through interaction with ITPR1 catalytic activity:S-adenosyl-L-homocysteine + H(2)O = L-homocysteine + adenosine.,cofactor:Binds 1 NAD per subunit.,cofactor:NAD.,pathway:Amino-acid biosynthesis; homocysteine biosynthesis; L-homocysteine from S-adenosyl-L-homocysteine: step 1/1.,similarity:Belongs to the adenosylhomocysteinase family., The protein encoded by this gene acts as a homotetramer and may be involved in the conversion of S-adenosyl-L-homocysteine to L-homocysteine and adenosine. Several transcript variants encoding different isoforms have been found for this



UpingBio technology Co.,Ltd


C Tel: 400-999-8863 ■ Email:UpingBio@163.com



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from Hela cells, primary antibody was diluted at 1:1000, 4°over night