



## VIP2 rabbit pAb

<b>Catalog No</b>	YP-Ab-11979
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human; Mouse
<b>Applications</b>	WB
<b>Gene Name</b>	PPIP5K2 HISPPD1 KIAA0433 VIP2
<b>Protein Name</b>	VIP2
<b>Immunogen</b>	Synthesized peptide derived from human VIP2 AA range: 266-316
<b>Specificity</b>	This antibody detects endogenous levels of VIP2 at Human/Mouse
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
<b>Dilution</b>	WB 1: 500-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	
<b>Cell Pathway</b>	Cytoplasm, cytosol .
<b>Tissue Specificity</b>	
<b>Function</b>	catalytic activity:ATP + 1D-myo-inositol 1,3,4,5,6-pentakisphosphate = ADP + diphospho-1D-myo-inositol tetrakisphosphate (isomeric configuration unknown).,catalytic activity:ATP + 1D-myo-inositol 5-diphosphate pentakisphosphate = ADP + 1D-myo-inositol bisdiphosphate tetrakisphosphate (isomeric configuration unknown).,catalytic activity:ATP + 1D-myo-inositol hexakisphosphate = ADP + 5-diphospho-1D-myo-inositol (1,2,3,4,6)pentakisphosphate.,caution:Although related to histidine acid phosphatases, it lacks the conserved active sites, suggesting that it has no phosphatase activity.,function:Bifunctional inositol kinase that catalyzes the formation of diphosphoinositol pentakisphosphate (InsP7 or PP-InsP5) and bi-diphosphoinositol tetrakisphosphate (InsP8 or PP2-InsP4). Converts inositol hexakisphosphate (InsP6) to InsP7. Also able to convert InsP7 to InsP8. Probably specifically mediates
<b>Background</b>	This gene encodes a member of the histidine acid phosphatase family of proteins. Despite containing a histidine acid phosphatase domain, the encoded protein



functions as an inositol pyrophosphate kinase, and is thought to lack phosphatase activity. This kinase activity is the mechanism by which the encoded protein synthesizes high-energy inositol pyrophosphates, which act as signaling molecules that regulate cellular homeostasis and other processes. This gene may be associated with autism spectrum disorder in human patients. [provided by RefSeq, Sep 2016],


**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Western blot analysis of lysates from THP-1 cells, primary antibody was diluted at 1:1000, 4° over night