







## PCGF5 rabbit pAb


Catalog No	YP-Ab-12033
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB;ELISA;IHC
Gene Name	PCGF5 RNF159
Protein Name	PCGF5
Immunogen	Synthesized peptide derived from human PCGF5 AA range: 65-115
Specificity	This antibody detects endogenous levels of PCGF5 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Storage Stability Synonyms	-20°C/1 year
	-20°C/1 year
Synonyms	-20°C/1 year  Nucleus . Nucleus, nucleoplasm . Recruited by the non-coding RNA Xist to specific nuclear foci that probably correspond to the inactivated X chromosome
Synonyms Observed Band	Nucleus . Nucleus, nucleoplasm . Recruited by the non-coding RNA Xist to
Synonyms Observed Band Cell Pathway	Nucleus . Nucleus, nucleoplasm . Recruited by the non-coding RNA Xist to
Synonyms  Observed Band  Cell Pathway  Tissue Specificity	Nucleus . Nucleus, nucleoplasm . Recruited by the non-coding RNA Xist to specific nuclear foci that probably correspond to the inactivated X chromosome function:Probable component of some Polycomb group (PcG) multiprotein complex, a complex required to maintain the transcriptionally repressive state of
Synonyms  Observed Band  Cell Pathway  Tissue Specificity  Function	Nucleus . Nucleus, nucleoplasm . Recruited by the non-coding RNA Xist to specific nuclear foci that probably correspond to the inactivated X chromosome function:Probable component of some Polycomb group (PcG) multiprotein complex, a complex required to maintain the transcriptionally repressive state of
Synonyms  Observed Band  Cell Pathway  Tissue Specificity  Function  Background  matters needing	Nucleus . Nucleus, nucleoplasm . Recruited by the non-coding RNA Xist to specific nuclear foci that probably correspond to the inactivated X chromosome function:Probable component of some Polycomb group (PcG) multiprotein complex, a complex required to maintain the transcriptionally repressive state of some genes.,similarity:Contains 1 RING-type zinc finger.,








## **Products Images**



Western blot analysis of lysates from MCF-7 cells, primary antibody was diluted at 1:1000, 4°over night



Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).