


MFS12 rabbit pAb


Catalog No	YP-Ab-12170
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB;ELISA;IHC
Gene Name	MFSD12 C19orf28
Protein Name	MFS12
Immunogen	Synthesized peptide derived from human MFS12 AA range: 78-128
Specificity	This antibody detects endogenous levels of MFS12 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Melanosome membrane ; Multi-pass membrane protein . Lysosome membrane ; Multi-pass membrane protein .
Cell Pathway Tissue Specificity	
-	Multi-pass membrane protein . Widely expressed, with high expression in primary melanocytes.
Tissue Specificity	Multi-pass membrane protein . Widely expressed, with high expression in primary melanocytes. caution:The sequence shown here is derived from an Ensembl automatic analysis pipeline and should be considered as preliminary data.,similarity:Belongs to the
Tissue Specificity Function	Multi-pass membrane protein . Widely expressed, with high expression in primary melanocytes. caution:The sequence shown here is derived from an Ensembl automatic analysis pipeline and should be considered as preliminary data.,similarity:Belongs to the
Tissue Specificity Function Background matters needing	Multi-pass membrane protein . Widely expressed, with high expression in primary melanocytes. caution:The sequence shown here is derived from an Ensembl automatic analysis pipeline and should be considered as preliminary data.,similarity:Belongs to the major facilitator superfamily.,



Products Images



Western blot analysis of lysates from HEK293 cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human Squamous cell carcinoma of lung. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).