



K1683 rabbit pAb

Catalog No	YP-Ab-12209
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;IHC
Gene Name	KIAA1683
Protein Name	K1683
Immunogen	Synthesized peptide derived from human K1683 AA range: 328-378
Specificity	This antibody detects endogenous levels of K1683 at Human
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Storage Stability Synonyms	-20°C/1 year
	-20°C/1 year
Synonyms	-20°C/1 year nucleus,mitochondrion,
Synonyms Observed Band	
Synonyms Observed Band Cell Pathway	
Synonyms Observed Band Cell Pathway Tissue Specificity	nucleus,mitochondrion, PTM:Phosphorvlated upon DNA damage, probably by ATM or ATRsequence
Synonyms Observed Band Cell Pathway Tissue Specificity Function	nucleus,mitochondrion, PTM:Phosphorvlated upon DNA damage, probably by ATM or ATRsequence
Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	nucleus,mitochondrion, PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,sequence caution:Wrong choice of frame.,similarity:Contains 6 IQ domains.,

Products Images



Western blot analysis of lysates from DU145 cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human Squamous cell carcinoma of lung. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).