



TRI11 rabbit pAb

Catalog No	YP-Ab-12265
Isotype	IgG
Reactivity	Human; Mouse; Rat
Applications	WB
Gene Name	TRIM11 RNF92
Protein Name	TRI11
Immunogen	Synthesized peptide derived from human TRI11 AA range: 128-178
Specificity	This antibody detects endogenous levels of TRI11 at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit, IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Cytoplasm. Nucleus.
Tissue Specificity	Ubiquitous.
Function	domain: The coiled-coil domain and the B30.2 domain are both necessary for interaction with humanin., function: May have E3 ubiquitin ligase activity. May contribute to the regulation of the intracellular level of humanin or humanin-containing proteins through the proteasomal degradation pathway., similarity: Belongs to the TRIM/RBCC family., similarity: Contains 1 B box-type zinc finger., similarity: Contains 1 B30.2/SPRY domain., similarity: Contains 1 RING-type zinc finger., subunit: Binds cytoplasmic tail of integrin alpha-1. Interacts with the humanin peptide., tissue specificity: Ubiquitous.,
Background	The protein encoded by this gene is a member of the tripartite motif (TRIM) family. The TRIM motif includes three zinc-binding domains, a RING, a B-box type 1 and a B-box type 2, and a coiled-coil region. This protein localizes to the nucleus and the cytoplasm. Its function has not been identified. [provided by RefSeq, Jul 2008],

**matters needing attention**

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

