



## MLIP rabbit pAb

Catalog No	YP-Ab-12266
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB;ELISA;IHC
Gene Name	MLIP C6orf142
Protein Name	MLIP
Immunogen	Synthesized peptide derived from human MLIP AA range: 216-266
Specificity	This antibody detects endogenous levels of MLIP at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000
Concentration	1 mg/ml
Concentration Purity	1 mg/ml ≥90%
	-
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90% -20°C/1 year
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year  Nucleus . Nucleus envelope . Nucleus, PML body . Cell membrane, sarcolemma ;
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90%  -20°C/1 year  Nucleus . Nucleus envelope . Nucleus, PML body . Cell membrane, sarcolemma ; Peripheral membrane protein ; Cytoplasmic side .
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90%  -20°C/1 year  Nucleus . Nucleus envelope . Nucleus, PML body . Cell membrane, sarcolemma ; Peripheral membrane protein ; Cytoplasmic side .
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90%  -20°C/1 year  Nucleus . Nucleus envelope . Nucleus, PML body . Cell membrane, sarcolemma ; Peripheral membrane protein ; Cytoplasmic side .
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	≥90%  -20°C/1 year  Nucleus . Nucleus envelope . Nucleus, PML body . Cell membrane, sarcolemma ; Peripheral membrane protein ; Cytoplasmic side .  Primarily expressed in heart and skeletal muscle. Also detected in liver.

## **Products Images**



Western blot analysis of lysates from 293T cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human uterus. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).