

## Amphiphysin II Monoclonal Antibody

matters needing attention	Avoid repeated freezing and thawing!
Background	
Function	
Tissue Specificity	Ubiquitous. Highest expression in the brain and muscle (PubMed:9182667). Expressed in oligodendrocytes (PubMed:27488240). Isoform IIA is expressed only in the brain, where it is detected in the gray matter, but not in the white matter (PubMed:27488240). Isoform BIN1 is widely expressed with highest expression in skeletal muscle.
Cell Pathway	[Isoform BIN1]: Nucleus . Cytoplasm . Endosome . Cell membrane, sarcolemma, T-tubule .; [Isoform IIA]: Cytoplasm .
Observed Band	11367DA
Synonyms	BIN1; AMPHL; Myc box-dependent-interacting protein 1; Amphiphysin II; Amphiphysin-like protein; Box-dependent myc-interacting protein 1; Bridging integrator 1
Storage Stability	-20°C/1 year
Purity	≥90%
Concentration	1 mg/ml
Dilution	WB 1:500-1:2000
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Source	Monoclonal, Mouse,IgG
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Specificity	Amphiphysin II Monoclonal Antibody detects endogenous levels of Amphiphysin II protein.
Immunogen	The antiserum was produced against synthesized peptide derived from the C-terminal region of human BIN1. AA range:521-570
Protein Name	Myc box-dependent-interacting protein 1
Gene Name	BIN1
Applications	WB
Reactivity	Human;Mouse;Rat
Isotype	IgG
Catalog No	YP-mAb-00767



## UpingBio technology Co.,Ltd





**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**