



β -1,4-GalNAc-T Monoclonal Antibody

Catalog No	YP-mAb-04290
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	CSGALNACT1
Protein Name	Chondroitin sulfate N-acetylgalactosaminyltransferase 1
Immunogen	The antiserum was produced against synthesized peptide derived from human CSGALNACT1. AA range:201-250
Specificity	β -1,4-GalNAc-T Monoclonal Antibody detects endogenous levels of β -1,4-GalNAc-T protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	$\geq 90\%$
Storage Stability	-20°C/1 year
Synonyms	CSGALNACT1; CHGN; GALNACT1; Chondroitin sulfate N-acetylgalactosaminyltransferase 1; CsGalNAcT-1; Chondroitin beta-1; 4-N-acetylgalactosaminyltransferase 1; Beta4GalNAcT-1
Observed Band	60kD
Cell Pathway	Golgi apparatus, Golgi stack membrane ; Single-pass type II membrane protein .
Tissue Specificity	Ubiquitous, with the highest levels in placenta, thyroid, bladder, prostate and adrenal gland. Detected at low levels in the other tissues examined.
Function	catalytic activity:UDP-N-acetyl-D-galactosamine + beta-D-glucuronyl-(1->3)-D-galactosyl-proteoglycan = UDP + N-acetyl-D-galactosaminyl-(1->4)-beta-D-glucuronyl-(1->3)-beta-D-galactosylproteoglycan.;function:Transfers 1,4-N-acetylgalactosamine (GalNAc) from UDP-GalNAc to the non-reducing end of glucuronic acid (GlcUA). Required for addition of the first GalNAc to the core tetrasaccharide linker and for elongation of chondroitin chains. Important role in chondroitin chain biosynthesis in cartilage.,online information:Chondroitin beta-1,4-N-acetylgalactosaminyltransferase 1,online information:GlycoGene database,PTM:N-glycosylated.,similarity:Belongs to the chondroitin N-acetylgalactosaminyltransferase family.,tissue specificity:Ubiquitous, with the highest levels in placenta, thyroid, bladder, prostate and adrenal gland. Detected



at low levels in the other tissues examined.,

Background

catalytic activity:UDP-N-acetyl-D-galactosamine + beta-D-glucuronyl-(1->3)-D-galactosyl-proteoglycan = UDP + N-acetyl-D-galactosaminyl-(1->4)-beta-D-glucuronyl-(1->3)-beta-D-galactosylproteoglycan.,function:Transfers 1,4-N-acetylgalactosamine (GalNAc) from UDP-GalNAc to the non-reducing end of glucuronic acid (GlcUA). Required for addition of the first GalNAc to the core tetrasaccharide linker and for elongation of chondroitin chains. Important role in chondroitin chain biosynthesis in cartilage.,online information:Chondroitin beta-1,4-N-acetylgalactosaminyltransferase 1,online information:GlycoGene database,PTM:N-glycosylated.,similarity:Belongs to the chondroitin N-acetylgalactosaminyltransferase family.,tissue specificity:Ubiquitous, with the highest levels in placenta, thyroid, bladder, prostate and adrenal gland. Detected at low levels in the other tissues examined.,

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images