






## FGFR1 Monoclonal Antibody

matters needing attention	Avoid repeated freezing and thawing!
Background	
Function	
Tissue Specificity	Detected in astrocytoma, neuroblastoma and adrenal cortex cell lines. Some isoforms are detected in foreskin fibroblast cell lines, however isoform 17, isoform 18 and isoform 19 are not detected in these cells.
Cell Pathway	Cell membrane; Single-pass type I membrane protein. Nucleus. Cytoplasm, cytosol. Cytoplasmic vesicle. After ligand binding, both receptor and ligand are rapidly internalized. Can translocate to the nucleus after internalization, or by translocation from the endoplasmic reticulum or Golgi apparatus to the cytosol, and from there to the nucleus.
Observed Band	55716DA
Synonyms	
Storage Stability	-20°C/1 year
Purity	≥90%
Concentration	1 mg/ml
Dilution	WB 1:500-1:2000
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Source	Monoclonal, Mouse,IgG
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Specificity	FGFR1 Monoclonal Antibody detects endogenous levels of protein.
Immunogen	Synthesized peptide derived from human protein . at AA range: 600-680
Protein Name	Fibroblast growth factor receptor 1 (FGFR-1) (EC 2.7.10.1) (Basic fibroblast growth factor receptor 1) (BFGFR) (bFGF-R-1) (Fms-like tyrosine kinase 2) (FLT-2) (N-sam) (Proto-oncogene c-Fgr) (CD antigen CD331)
Gene Name	FGFR1 BFGFR CEK FGFBR FLG FLT2 HBGFR
Applications	WB
Reactivity	Human;Mouse;Rat
Isotype	IgG
Catalog No	YP-mAb-04901



## UpingBio technology Co.,Ltd





**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**