





2AAA Monoclonal Antibody

Catalog No	YP-mAb-04967
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	PPP2R1A
Protein Name	Serine/threonine-protein phosphatase 2A 65 kDa regulatory subunit A alpha isoform (Medium tumor antigen-associated 61 kDa protein) (PP2A subunit A isoform PR65-alpha) (PP2A subunit A isoform R1-alpha)
Immunogen	Synthesized peptide derived from human protein . at AA range: 500-580
Specificity	2AAA Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	64kD
Cell Pathway	Cytoplasm . Nucleus . Chromosome, centromere . Lateral cell membrane . Cell projection, dendrite . Centromeric localization requires the presence of BUB1
Tissue Specificity	Brain, Cajal-Retzius cell, Colon, Placenta, Testis,
Function	domain:Each HEAT repeat appears to consist of two alpha helices joined by a hydrophilic region, the intrarepeat loop. The repeat units may be arranged laterally to form a rod-like structure.,function:The PR65 subunit of protein phosphatase 2A serves as a scaffolding molecule to coordinate the assembly of the catalytic subunit and a variable regulatory B subunit.,similarity:Belongs to the phosphatase 2A regulatory subunit A family.,similarity:Contains 15 HEAT repeats.,subunit:PP2A consists of a common heterodimeric core enzyme, composed of a 36 kDa catalytic subunit (subunit C) and a 65 kDa constant regulatory subunit (PR65 or subunit A), that associates with a variety of regulatory subunits. Proteins that associate with the core dimer include three families of regulatory subunits B (the R2/B/PR55/B55, R3/B"/PR72/PR130/PR59 and R5/B'/B56 families), the 48 kDa variable regulatory subunit,



UpingBio technology Co.,Ltd







implicated in the negative control of cell growth and division. It consists of a common heteromeric core enzyme, which is composed of a catalytic subunit and a constant regulatory subunit, that associates with a variety of regulatory subunits. The constant regulatory subunit A serves as a scaffolding molecule to coordinate the assembly of the catalytic subunit and a variable regulatory is the other transfer of the catalytic subunit and a variable regulatory is the other transfer of the catalytic subunit and a variable regulatory is the other transfer of the catalytic subunit and a variable regulatory is the other transfer of the catalytic subunit and a variable regulatory is the other transfer of the catalytic subunit and a variable regulatory is the other transfer of the catalytic subunit and a variable regulatory is the other transfer of the catalytic subunit and a variable regulatory is the other transfer of the catalytic subunit and a variable regulatory is the other transfer of the catalytic subunit and a variable regulatory is the other transfer of the catalytic subunit and a variable regulatory is the other transfer of the catalytic subunit and the catalytic subunit and the catalytic subun gene encodés an alpha isoform of the constant regulatory subunit A. Alternatively spliced transcript variants have been described. [provided by RefSeq, Apr 2010],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images