





RBPJL Monoclonal Antibody

| Catalog No | YP-mAb-05029 |
|---------------------------|---|
| Isotype | IgG |
| Reactivity | Human;Mouse |
| Applications | WB |
| Gene Name | RBPJL RBPL RBPSUHL |
| Protein Name | Recombining binding protein suppressor of hairless-like protein (Transcription factor RBP-L) |
| Immunogen | Synthesized peptide derived from human protein . at AA range: 140-220 |
| Specificity | RBPJL Monoclonal Antibody detects endogenous levels of protein. |
| Formulation | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. |
| Source | Monoclonal, Mouse,IgG |
| Purification | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-1:2000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | |
| Observed Band | 56kD |
| Cell Pathway | Nucleus . |
| Tissue Specificity | |
| Function | function:Putative transcription factor, which cooperates with EBNA2 to activate transcription.,similarity:Belongs to the Su(H) family.,similarity:Contains 1 IPT/TIG domain.,subunit:Interacts weakly with EBNA2. Does not interact with any Notch proteins., |
| Background | This gene encodes a member of the suppressor of hairless protein family. A similar protein in mouse is a transcription factor that binds to DNA sequences almost identical to that bound by the Notch receptor signaling pathway transcription factor recombining binding protein J. The mouse protein has been shown to activate transcription in concert with Epstein-Barr virus nuclear antigen-2. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul 2013], |
| matters needing attention | Avoid repeated freezing and thawing! |
| | |



UpingBio technology Co.,Ltd





Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images