



# IFIT3 Monoclonal Antibody

<b>Catalog No</b>	YP-mAb-05087
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse;
<b>Applications</b>	WB
<b>Gene Name</b>	IFIT3 CIG-49 IFI60 IFIT4 ISG60
<b>Protein Name</b>	Interferon-induced protein with tetratricopeptide repeats 3 (IFIT-3) (CIG49) (ISG-60) (Interferon-induced 60 kDa protein) (IFI-60K) (Interferon-induced protein with tetratricopeptide repeats 4) (IFIT-
<b>Immunogen</b>	Synthesized peptide derived from human protein . at AA range: 330-410
<b>Specificity</b>	IFIT3 Monoclonal Antibody detects endogenous levels of protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse,IgG
<b>Purification</b>	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-1:2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	53kD
<b>Cell Pathway</b>	Cytoplasm . Mitochondrion .
<b>Tissue Specificity</b>	Expression significantly higher in peripheral blood mononuclear cells (PBMCs) and monocytes from systemic lupus erythematosus (SLE) patients than in those from healthy individuals (at protein level). Spleen, lung, leukocytes, lymph nodes, placenta, bone marrow and fetal liver.
<b>Function</b>	similarity:Belongs to the IFIT family.,similarity:Contains 8 TPR repeats.,
<b>Background</b>	similarity:Belongs to the IFIT family.,similarity:Contains 8 TPR repeats.,
<b>matters needing attention</b>	Avoid repeated freezing and thawing!
<b>Usage suggestions</b>	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



## Products Images