



# CAN6 Monoclonal Antibody

<b>Catalog No</b>	YP-mAb-05413
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse
<b>Applications</b>	WB
<b>Gene Name</b>	CAPN6 CALPM CANPX
<b>Protein Name</b>	Calpain-6 (Calpain-like protease X-linked) (Calpamodulin) (CalpM)
<b>Immunogen</b>	Synthesized peptide derived from part region of human protein
<b>Specificity</b>	CAN6 Monoclonal Antibody detects endogenous levels of protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse,IgG
<b>Purification</b>	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-1:2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	70kD
<b>Cell Pathway</b>	Cytoplasm, perinuclear region . Cytoplasm, cytoskeleton, spindle . During mitose associated with the mitotic spindle. At telophase colocalized to the midbody spindle.
<b>Tissue Specificity</b>	Expressed only in placenta.
<b>Function</b>	function:Not known; does not seem to have protease activity as it has lost the active site residues.,similarity:Belongs to the peptidase C2 family.,similarity:Contains 1 C2 domain.,similarity:Contains 1 calpain catalytic domain.,tissue specificity:Expressed only in placenta.,
<b>Background</b>	Calpains are ubiquitous, well-conserved family of calcium-dependent, cysteine proteases. The calpain proteins are heterodimers consisting of an invariant small subunit and variable large subunits. The large subunit possesses a cysteine protease domain, and both subunits possess calcium-binding domains. Calpains have been implicated in neurodegenerative processes, as their activation can be triggered by calcium influx and oxidative stress. The protein encoded by this gene is highly expressed in the placenta. Its C-terminal region lacks any homology to the calmodulin-like domain of other calpains. The protein lacks critical active site residues and thus is suggested to be proteolytically inactive. The protein may play a role in tumor formation by inhibiting apoptosis and promoting angiogenesis.



[provided by RefSeq, Nov 2009],

**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**