



# CCDC8 Monoclonal Antibody

<b>Catalog No</b>	YP-mAb-05485
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse;
<b>Applications</b>	WB
<b>Gene Name</b>	CCDC8
<b>Protein Name</b>	Coiled-coil domain-containing protein 8
<b>Immunogen</b>	Synthesized peptide derived from part region of human protein
<b>Specificity</b>	CCDC8 Monoclonal Antibody detects endogenous levels of protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse,IgG
<b>Purification</b>	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-1:2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	59kD
<b>Cell Pathway</b>	Cytoplasm . Cytoplasm, cytoskeleton, microtubule organizing center, centrosome
<b>Tissue Specificity</b>	Widely expressed with low levels in spleen, skeletal muscle, small intestine, kidney and liver.
<b>Function</b>	PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,
<b>Background</b>	This gene encodes a coiled-coil domain-containing protein. The encoded protein functions as a cofactor required for p53-mediated apoptosis following DNA damage, and may also play a role in growth through interactions with the cytoskeletal adaptor protein obscurin-like 1. Mutations in this gene are a cause of 3M syndrome-3 (3M3). [provided by RefSeq, Dec 2011],
<b>matters needing attention</b>	Avoid repeated freezing and thawing!
<b>Usage suggestions</b>	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



## Products Images