



DNA2 Monoclonal Antibody

Catalog No	YP-mAb-05526
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	DNA2 DNA2L KIAA0083
Protein Name	DNA replication ATP-dependent helicase/nuclease DNA2 (hDNA2) (DNA replication ATP-dependent helicase-like homolog) [Includes: DNA replication nuclease DNA2 (EC 3.1.-.-); DNA replication ATP-dependent
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	DNA2 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	116kD
Cell Pathway	Nucleus. Mitochondrion. Was initially reported to be exclusively mitochondrial (PubMed:18995831). However, it was later shown to localize both in mitochondrion and nucleus (PubMed:19487465). .
Tissue Specificity	Bone marrow,Colon,Duodenum,Lymph,
Function	function:May function in chromosomal DNA replication.,similarity:Belongs to the DNA2/NAM7 helicase family.,
Background	This gene encodes a member of the DNA2/NAM7 helicase family. The encoded protein is a conserved helicase/nuclease involved in the maintenance of mitochondrial and nuclear DNA stability. Mutations in this gene are associated with autosomal dominant progressive external ophthalmoplegia-6 (PEOA6) and Seckel syndrome 8. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Sep 2014],
matters needing attention	Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images