





ACSL5 Monoclonal Antibody

Catalog No	YP-mAb-05702
Isotype	IgG
Reactivity	Human;Rat;Mouse
Applications	WB
Gene Name	ACSL5 ACS5 FACL5 UNQ633/PRO1250
Protein Name	Long-chain-fatty-acidCoA ligase 5 (EC 6.2.1.3) (Long-chain acyl-CoA synthetase 5) (LACS 5)
Immunogen	Synthesized peptide derived from human protein . at AA range: 460-540
Specificity	ACSL5 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	75kD
Cell Pathway	Mitochondrion . Endoplasmic reticulum . Mitochondrion outer membrane ; Single-pass type III membrane protein . Endoplasmic reticulum membrane ; Single-pass type III membrane protein . Cell membrane .
Tissue Specificity	Colon,
Function	catalytic activity:ATP + a long-chain carboxylic acid + CoA = AMP + diphosphate + an acyl-CoA.,cofactor:Magnesium.,function:Activation of long-chain fatty acids for both synthesis of cellular lipids, and degradation via beta-oxidation. Utilizes a wide range of saturated fatty acids with a preference for C16-C18 unsaturated fatty acids.,similarity:Belongs to the ATP-dependent AMP-binding enzyme family.,
Background	The protein encoded by this gene is an isozyme of the long-chain fatty-acid-coenzyme A ligase family. Although differing in substrate specificity, subcellular localization, and tissue distribution, all isozymes of this family convert free long-chain fatty acids into fatty acyl-CoA esters, and thereby play a key role in lipid biosynthesis and fatty acid degradation. This isozyme is highly expressed in uterus and spleen, and in trace amounts in normal brain, but has markedly increased levels in malignant gliomas. This gene functions in mediating fatty acid-induced glioma cell growth. Three transcript variants encoding two different isoforms have been found for this gene. [provided by RefSeq, Jul 2008],



UpingBio technology Co.,Ltd





matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images