



NOG2 Monoclonal Antibody

Catalog No	YP-mAb-05879
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	GNL2 NGP1
Protein Name	Nucleolar GTP-binding protein 2 (Autoantigen NGP-1)
Immunogen	Synthesized peptide derived from human protein . at AA range: 670-750
Specificity	NOG2 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	80kD
Cell Pathway	Nucleus, nucleolus .
Tissue Specificity	Widely expressed, with the highest expression level in testis.
Function	function:GTPase that associates with pre-60S ribosomal subunits in the nucleolus and is required for their nuclear export and maturation.,similarity:Belongs to the MMR1/HSR1 GTP-binding protein family. NOG2 subfamily.,tissue specificity:Ubiquitously expressed at relatively low levels in all human tissues tested, with the highest level of expression in the testes.,
Background	function:GTPase that associates with pre-60S ribosomal subunits in the nucleolus and is required for their nuclear export and maturation.,similarity:Belongs to the MMR1/HSR1 GTP-binding protein family. NOG2 subfamily.,tissue specificity:Ubiquitously expressed at relatively low levels in all human tissues tested, with the highest level of expression in the testes.,
matters needing attention	Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images