



PLXB1 Monoclonal Antibody

Catalog No	YP-mAb-05941
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	PLXNB1 KIAA0407 PLXN5 SEP
Protein Name	Plexin-B1 (Semaphorin receptor SEP)
Immunogen	Synthesized peptide derived from human protein . at AA range: 90-170
Specificity	PLXB1 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	234kD
Cell Pathway	[Isoform 1]: Cell membrane ; Single-pass type I membrane protein .; [Isoform 2]: Secreted .; [Isoform 3]: Secreted .
Tissue Specificity	Highly expressed in fetal kidney, and at slightly lower levels in fetal brain, lung and liver.
Function	disease:Overexpressed and constitutively tyrosine phosphorylated in colon, liver, pancreas and gastric carcinoma cell lines. Overexpression increases MET activation and promotes invasive growth.,function:Receptor for SEMA4D. Plays a role in RHOA activation and subsequent changes of the actin cytoskeleton. Plays a role in axon guidance, invasive growth and cell migration.,PTM:Phosphorylated on tyrosine residues by ERBB2 and MET upon SEMA4D binding.,PTM:Proteolytic processing favors heterodimerization with PLXNB2 and SEMA4D binding.,similarity:Belongs to the plexin family.,similarity:Contains 1 Sema domain.,similarity:Contains 3 IPT/TIG domains.,subunit:Monomer, and heterodimer with PLXNB2 after proteolytic processing. Binds RAC1 that has been activated by GTP binding. Interaction with SEMA4D promotes binding of cytoplasmic ligands. Binds PLXNA1 (By similarity). Binds ARHGEF11, ARHGEF12, E

**Background**

disease:Overexpressed and constitutively tyrosine phosphorylated in colon, liver, pancreas and gastric carcinoma cell lines. Overexpression increases MET activation and promotes invasive growth.,function:Receptor for SEMA4D. Plays a role in RHOA activation and subsequent changes of the actin cytoskeleton. Plays a role in axon guidance, invasive growth and cell migration.,PTM:Phosphorylated on tyrosine residues by ERBB2 and MET upon SEMA4D binding.,PTM:Proteolytic processing favors heterodimerization with PLXNB2 and SEMA4D binding.,similarity:Belongs to the plexin family.,similarity:Contains 1 Sema domain.,similarity:Contains 3 IPT/TIG domains.,subunit:Monomer, and heterodimer with PLXNB2 after proteolytic processing. Binds RAC1 that has been activated by GTP binding. Interaction with SEMA4D promotes binding of cytoplasmic ligands. Binds PLXNA1 (By similarity). Binds ARHGEF11, ARHGEF12, ERBB2, MET, MST1R, RND1, NRP1 and NRP2.,tissue specificity:Highly expressed in fetal kidney, and at slightly lower levels in fetal brain, lung and liver.,

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images