



RPGEF6 Monoclonal Antibody

Catalog No	YP-mAb-06020
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	RAPGEF6 PDZGEF2
Protein Name	Rap guanine nucleotide exchange factor 6 (PDZ domain-containing guanine nucleotide exchange factor 2) (PDZ-GEF2) (RA-GEF-2)
Immunogen	Synthesized peptide derived from human protein . at AA range: 460-540
Specificity	RPGEF6 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	176kD
Cell Pathway	Cytoplasm . Cell membrane . Upon binding to M-Ras, it translocates to the plasma membrane.
Tissue Specificity	Isoform 3 has highest expression levels in the brain, heart, liver, lung and placenta and is barely detectable in skeletal muscle, kidney and pancreas.
Function	function:Guanine nucleotide exchange factor (GEF) for Rap1A, Rap2A and M-Ras GTPases. Does not interact with cAMP.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 cyclic nucleotide-binding domain.,similarity:Contains 1 N-terminal Ras-GEF domain.,similarity:Contains 1 PDZ (DHR) domain.,similarity:Contains 1 Ras-associating domain.,similarity:Contains 1 Ras-GEF domain.,subcellular location:Upon binding to M-Ras, it translocates to the plasma membrane.,subunit:Interacts with the second PDZ domain of human PTP1e.,tissue specificity:Isoform 3 has highest expression levels in the brain, heart, liver, lung and placenta and is barely detectable in skeletal muscle, kidney and pancreas.,
Background	function:Guanine nucleotide exchange factor (GEF) for Rap1A, Rap2A and M-Ras GTPases. Does not interact with cAMP.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 cyclic nucleotide-binding domain.,similarity:Contains 1 N-terminal Ras-GEF domain.,similarity:Contains 1



PDZ (DHR) domain.,similarity:Contains 1 Ras-associating domain.,similarity:Contains 1 Ras-GEF domain.,subcellular location:Upon binding to M-Ras, it translocates to the plasma membrane.,subunit:Interacts with the second PDZ domain of human PTP1e.,tissue specificity:Isoform 3 has highest expression levels in the brain, heart, liver, lung and placenta and is barely detectable in skeletal muscle, kidney and pancreas.,

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images